TECHNICAL INFORMATION BULLETIN

FIFRA Section 2(ee) Recommendation

This recommendation is made as permitted by Section 2(ee) of FIFRA, as amended, and has not been submitted to or approved by the U. S. Environmental Protection Agency.

FOR DISTRIBUTION AND USE WITHIN THE STATES OF TEXAS ONLY.

Sonata[®] EPA Reg. No. 69592-13

Reduced Sonata application rates recommended, for the control of Powdery Mildew only, when Sonata is used in a tank mix with a systemic fungicide registered for use on the crops and diseases specified in the accompanying application table.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

This FIFRA Section 2(ee) recommendation contains new or additional directions for use of this product, which may not appear on the package label. Read and carefully observe the applicable directions, restrictions and precautionary statements plus all the other information appearing on the EPA-registered product label.

Sonata ASO may be tank-mixed with other registered fungicides to enhance plant disease control. Sonata ASO cannot be mixed with any product with prohibition against such mixing. When tank-mixing Sonata ASO with other registered pesticides, always read and follow all use directions, restrictions, and precautions of both Sonata ASO and the tank-mix partner(s). Use of the resulting tank mix must be in accordance with the more restrictive label limitations and precautions. Do not exceed label dosage rates for the tank mix partner and follow the dosage rates for Sonata set forth in this recommendation.

This recommendation must be in the possession of the user at the time of pesticide application.



TECHNICAL INFORMATION BULLETIN

FIFRA Section 2(ee) Recommendation

FOR DISTRIBUTION AND USE WITHIN THE STATES OF TEXAS ONLY.

Sonata[®] EPA Reg. No. 69592-13

Reduced Sonata application rates recommended, for the control of Powdery Mildew only, when Sonata is used in a tank mix with a systemic fungicide registered for use on the crops and diseases specified in the following application table.

Crops	Diseases	Rate qt/acre	Application Instructions
Cucurbits Cucumber Cantaloupe Melon Muskmelon Squash Watermelon and other cucurbits	Powdery Mildew Erysiphe spp. Sphaerotheca spp.	1	Begin applications soon after emergence or transplant when environmental conditions and plant stage are conducive to disease development. Repeat on 7- to 14-day intervals or as needed. Use higher rates and short application intervals under heavy disease pressure. For improved performance, use Sonata ASO in a tank mix or rotational program with other registered fungicides.
Grape	Powdery Mildew Uncinula necator	1	Begin applications at prebloom. Continue at 7- to 14-day intervals or as needed. Use higher rates and shorter application intervals under heavy disease pressure. Use of a spreader/sticker or wetting agent is recommended to ensure optimum control. (Begin application when new shoots are ½ to 1½ inches long. Repeat when shoots are 3 to 5 inches long, when shoots are 8 to 10 inches long and then at 7- to 14-day intervals until disease conditions no longer exist.)
Pome Fruit Apple Crabapple Pear Quince Mayhaw and other pome fruit crops	Powdery Mildew Podosphaera leucotricha	1	Begin application at tight cluster, or sooner, if conditions are conducive to disease development. Repeat applications through the second cover spray at 7- to 14-day intervals. Additional sprays beyond second cover may be needed on susceptible varieties or when environmental conditions are conducive to rapid disease development or under heavy disease pressure. Use higher rates and shorter spray intervals when conditions are conducive to rapid disease development or heavy disease pressure.

Sonata[®] is a registered trademark of AgraQuest, Inc. [®] Copyright AgraQuest, Inc. 2012

Label Code: 050112 Page 2 0f 2

