



This seed has been treated with thiamethoxam insecticide and fludioxonil, mefenoxam, thiabendazole and sedaxane fungicides.

Do not use for feed, food, or oil purposes

Ground Water Advisory:

Thiamethoxam has properties and characteristics associated with chemicals detected in ground water. This chemical may leach into the ground water if used in areas where soils are permeable, particularly where the water table is shallow. Mefenoxam is known to leach through soil into ground water under certain conditions as a result of agricultural use. This chemical may leach into the ground water if used in areas where soils are permeable, particularly where the water table is shallow.

Pollinator Precautions:

Thiamethoxam is highly toxic to bees, and effects are possible as a result of exposure to translocated residues in blooming crops.

Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticide remain in the ethanol by-products that are used in agronomic practice.

Do not allow children, pets, or livestock to have access to treated seed.

Store away from feeds and foodstuffs.

Wear long-sleeved shirt, long pants and chemical resistant gloves when handling treated seed.

Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading.

Treated seed must be planted into the soil at a depth greater than 1 inch.

Dispose of all excess treated seed. Leftover treated seed may be doublesown around the headland or buried away from water sources in accordance with local requirements.

Do not contaminate water bodies when disposing of planting equipment wash waters.

Dispose of seed packaging in accordance with local requirements.

Do not graze or feed livestock on treated areas for 45 days after planting seed treated with THX/MXM/FLD/TBZ/SDX FS .

In the event of a crop failure or harvest of a crop grown from THX/MXM/FLD/TBZ/SDX FS treated seed, the field may be replanted according to the following schedule:

Immediate Plantback	Minimum 30-Day Plantback Interval	Minimum 120-Day Plantback Interval
Cereal Grains: Barley, Oats, Rye, Triticale, and Wheat Peas, Dried Shelled: Chickpea (Garbanzo Bean), Field Pea, Lentil, and Pigeon Pea Soybean	Alfalfa Brassica (Cole) Leafy Vegetables Cereal Grains (buckwheat, corn, pearl millet, proso, popcorn, rice (dry-seeded), sorghum, teosinte, wild rice Cotton Cucurbit Vegetables Fruiting Vegetables Globe Artichoke Leafy Vegetables Legume Vegetables (Succulent or Dried) Mint: Peppermint and Spearmint Oilseeds: Borage, Crambe, Flax Seed, Mustard Seed, and Safflower Peanut Root Vegetables Strawberry Sunflower Tobacco	All Other Crops

For any other crop, the minimum plant back interval is 120 days from the date the THX/MXM/FLD/TBZ/SDX FS treated seed was planted. A cover crop other than the crops listed above that is planted for erosion control or soil improvement may be planted sooner than the 120 day interval; however, the crop may not be grazed or harvested for food or feed.


Do not use at a rate that will result in more than 0.083 lb thiamethoxam (37.8 grams ai/A) per acre, 0.004 lb fludioxonil (2.0 grams ai/A) per acre and 0.013 lb mefenoxam (5.7 grams ai/A) per acre per calendar year as a seed treatment application.

This seed has been treated with 0.0756 mg ai thiamethoxam, 0.0113 mg mefenoxam, and 0.0038 mg fludioxonil per seed. x Do not apply a neonicotinoid insecticide within 45 days of planting seed treated with THX/MXM/FLD/TBZ/SDX.

7339 D0116



Loveland Products, Inc. • PO Box 1286 • Greeley, CO 80632



This seed has been treated with thiamethoxam insecticide and fludioxonil, mefenoxam, thiabendazole and sedaxane fungicides.

Do not use for feed, food, or oil purposes

Ground Water Advisory:

Thiamethoxam has properties and characteristics associated with chemicals detected in ground water. This chemical may leach into the ground water if used in areas where soils are permeable, particularly where the water table is shallow. Mefenoxam is known to leach through soil into ground water under certain conditions as a result of agricultural use. This chemical may leach into the ground water if used in areas where soils are permeable, particularly where the water table is shallow.

Pollinator Precautions:

Thiamethoxam is highly toxic to bees, and effects are possible as a result of exposure to translocated residues in blooming crops.

Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticide remain in the ethanol by-products that are used in agronomic practice.

Do not allow children, pets, or livestock to have access to treated seed.

Store away from feeds and foodstuffs.

Wear long-sleeved shirt, long pants and chemical resistant gloves when handling treated seed.

Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading.

Treated seed must be planted into the soil at a depth greater than 1 inch.

Dispose of all excess treated seed. Leftover treated seed may be doublesown around the headland or buried away from water sources in accordance with local requirements.

Do not contaminate water bodies when disposing of planting equipment wash waters.

Dispose of seed packaging in accordance with local requirements.

Do not graze or feed livestock on treated areas for 45 days after planting seed treated with THX/MXM/FLD/TBZ/SDX FS .

In the event of a crop failure or harvest of a crop grown from THX/MXM/FLD/TBZ/SDX FS treated seed, the field may be replanted according to the following schedule:

Immediate Plantback	Minimum 30-Day Plantback Interval	Minimum 120-Day Plantback Interval
Cereal Grains: Barley, Oats, Rye, Triticale, and Wheat Peas, Dried Shelled: Chickpea (Garbanzo Bean), Field Pea, Lentil, and Pigeon Pea Soybean	Alfalfa Brassica (Cole) Leafy Vegetables Cereal Grains (buckwheat, corn, pearl millet, proso, popcorn, rice (dry-seeded), sorghum, teosinte, wild rice Cotton Cucurbit Vegetables Fruiting Vegetables Globe Artichoke Leafy Vegetables Legume Vegetables (Succulent or Dried) Mint: Peppermint and Spearmint Oilseeds: Borage, Crambe, Flax Seed, Mustard Seed, and Safflower Peanut Root Vegetables Strawberry Sunflower Tobacco	All Other Crops

For any other crop, the minimum plant back interval is 120 days from the date the THX/MXM/FLD/TBZ/SDX FS treated seed was planted. A cover crop other than the crops listed above that is planted for erosion control or soil improvement may be planted sooner than the 120 day interval; however, the crop may not be grazed or harvested for food or feed.

Do not use at a rate that will result in more than 0.083 lb thiamethoxam (37.8 grams ai/A) per acre, 0.004 lb fludioxonil (2.0 grams ai/A) per acre and 0.013 lb mefenoxam (5.7 grams ai/A) per acre per calendar year as a seed treatment application.

This seed has been treated with 0.0756 mg ai thiamethoxam, 0.0113 mg mefenoxam, and 0.0038 mg fludioxonil per seed. x Do not apply a neonicotinoid insecticide within 45 days of planting seed treated with THX/MXM/FLD/TBZ/SDX.

7339 D0116



Loveland Products, Inc. • PO Box 1286 • Greeley, CO 80632