

ACTIVE INGREDIENT:

TICKS

Permethrin (3-phenoxyphenyl) methyl (+/-) cis, trans-3-(2,2-dichloroethenyl)-2,2dimethylcyclopropanecarboxylate*: 0.25%

cis/trans ratio: Min. 35% (+/-) cis, Max. 65% (+/-) trans

EPA Reg. No. 53883-33 EPA Est. No. 44616-MO-1^(A), EPA Est. No. 4-NY-1^(B) First letter of lot # corresponds to EPA Est # where product was produced.

KEEP OUT OF REACH OF CHILDREN

LICE

MITES

See back/side panels for additional precautionary statements

NET CONTENTS: 4 POUNDS

PASADENA, TEXAS 77507 **2903 GENOA-RED BLUFF** CONTROL SOLUTIONS, INC. Manufactured for

Dust lightly. Apply up to 8 times per season. May be applied until 1 day before harvest. Dust lightly, Apply up to 5 times per Season. May be applied until 14 days before arvest. Dust lightly. Do not apply after petal fall.

Navel Orangeworm, Codling Moth, Walnut Husk Fly Peach Twig Borer, Driental Fruit Moth, Green Fruitworm, Lesser Peachtree Borer, Plum Curculio, Tarnished Plant Bug

Uust lightly. Dust lightly. Ap<mark>pl</mark>y to the fem stage ot asparagus plant after speer harvest when adult stage of the asparagus beetle, lygus bugs and the adult stage of the . Dust nahtly. Apply up to 6 times per season. May be applied until day of harvest Dust lightly. Apply up to 6 times per season. May be applied until 1 day before harvest. ust lightly. Apply up to 8 times per season. May be applied until 14 days before harvest Just lightly. Apply up to 8 times per season. May be applied until 3 days before harvest. Just lightly. Apply up to 6 times per season. May be applied until 1 day before harvest. Dust lightly. Apply up to 10 times per season. May be applied until 3 days before harvest

Just lightly. Apply up to 5 times per season. May be applied until 1 day before harvest.

Dust lightly. Apply up to 5 times per season. May be applied until i day betore narvest.

Dust lightly. Apply up to 8 times per season. May be applied until 1 day before harvest.

Dust lightly. Apply up to 4 times per season. May be applied until 1 day before harvest.

Dust lightly. Apply up to 8 times per season. May be applied until harvest.

omato rruitworm, Cabbage Looper, Iomato Prinworm, Vegetable Leatminer, Beet Armyworm, Southern Armyworm, Colorado Potato I Corn Earworm, European Com Borer, Fall Armyworm, Southern Armyworm, Corn Rootworm (adult), Cutworm, Aster Leathopper, Flea Beetle olorado Potato Beetle, Potato Leafhopper, Aster Leafhopper, Potato Flea Beetle, Potato Aphid, Potato Tuberworm, Potato Psyllid, Cabbage Loop Pepper Weevil, Vegetable Leafminer, Cabbage Looper, Flea Beetle, European Corn Borer Dnion Thrips, Armyworm, Cutworm, Leatminer, Union Maggot (adult), Stink Bug Colorado Potato Beetle, Vegetable Leafminer, Cabbage Looper, Flea Beetle /egetable Leatminer, Cabbage Looper, Cutworm Complex, Beet Armyworm, Fall Armyworm, Corn Earworm, Leathopper, European Corn Borer

Cabbage Looper, Imported Cabbageworm, Diamondback Morth, Cabbage Aphid. Leatiminer, Squusa'n bug, Cabbage Looper, Cucumber Geetle, Galdit), Cutworm, Melon Worm, Pickle Worm, Plant Bug (including Lygus and Stink Bug), Rindworm, Squa

Sabbage Looper, Imported Cabbageworm, Diamondback Moth, Cabbage Aphid Cutworm Complex, Asparagus Beetle TREGET PEST

USE ON LIVESTOCK

hybrication is prohibited directly into sewers or drains, or to any area like a guider where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. Do may hybrication is prohibited directly into sewers or drains. As the supplication is prohibited directly into sewers or drains a milk lake list by prevent contamination from spray and from dead or falling insects. Remove or cover milking utensils before specifies. Observe milking utensils before any water or drains of run-off. Do not make applications during rain. Except when applying to pets, do not apply this product in a way that will contact workers or other persons either directly or through drift. Except when applying to pets, only protected handlers may be in the area during application. Do not breathe dusts, vapors, or spray mist. Do not enter or allow others to enter until dusts have settled.

Beet Armyworm, Cabbage Looper, Umnivorous Leatroller

ppanese Beetle, Asparagus Beetle, Lygus Bug

Jse of handheld power duster equipment is prohibited. t is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DIRECTIONS FOR USE

sacues

Asparagus Fern GR0P

тој јаамс

Peppers (Bell)

egegge

Asparagus

CROP

Dinons (Dry bulb and Garlic)

TREE CROPS: DIRECTIONS FOR APPLICATION Dust lightly, using a dust gun or rotary dus

DRNAMENTALS: DIRECTIONS FOR APPLICATION

сысту, геписе (пеаа апа теат) магяеу, кпираго, эртпасп

ucumber, Muskmelons, Pumpkin, Squash, Watermelon

Jset should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughlyand change into clean User should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. User should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. USER SAFETY RECOMMENDATION Follow manutacturer's instructions for deaning/ining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's byncentrate. Do not reuse them.

USER SAFETY REQUIREMENTS Applicators and other handlers must weer:
Long-sleeved shirt and brong pants, shoes plus socks, chemical-resistant gloves, and a NIOSH-approved respirator with: a dust/mist with MSHA/NIOSH approval number
Perfix TG-31 or any M, R, P, or HE filter. Application with motorized equipment is prohibited.

esistant to this product are listed below. If you want more options, follow the instructions for category E on an EPA chemical resistance РЕВЅОИАL РВОТЕСТІУЕ ЕQUIPMENT:

ENVIRONMENTAL HAZARDS

This besticals ear strengly takes to aquatic organisms, including flath and invertibrates. To protect the environment, do not allow pesticide for run off into storm drains, drainage disches waters will help ensure that whind or rain does not blow or wash disches the treatment area.

Pesticide off the treatment area.

PHYSICAL AND CHEMICAL HAZARDS

Have the product container or labels with you where calling besing control center or defects or defects or good treatment. For formation on this product, contact the National Breatiste Information or the product contact the National State (1906) 945-4508, Monday-Hiday, 230 PM PST, You may also contact the National Present (1906) 945-4508. day or night, for emergency made in teatment information or produced the National Psychiatric Psy ince person or incentar. If person is not beseiting call 911 or an ambulance; then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice. Move person to tresh air. bəledni t Call a poison control center or doctor tor treatment advice.

. Take off contaminated clothing. Binse skin immediately with plenty of water for 16-20 minytes. . Call a poison control center or doctor for treatment advice. GIA T2817

HAZANDS TO HELEMENTS.

HAZANDS TO HELEMENTS.

HAZANDS TO HELEMENTS.

HAZANDS TO HELEMENTS.

HAZANDS TO HUMANS AND DOMESTIC ANIMALS.

Charuld a veletrinarian before using this product on debullated, aged, pregnant or nursing animals on medication. Sensitivities may occur after using ANY pesticide Charuld a veletrinarian before using this product on debullated, aged, pregnant or nursing animals on medication. Sensitivity occur after using ANY pesticide product or debullativities may our pay with a mild seap and fines with large anounts of water. It signs consult a veletrinarian immediately be a pay east. Accidental application to cats and/or grooming a recently treated dog may result in tremors and/or uncoordinated muscle movements. If this occurs, immediately veletrinarian immediately veletrinary occurs, immediately occurs, immediately

PRECAUTIONARY STATEMENTS FOR PET USE - USE ONLY ON DOGS

LIMITATIONS DE LEAGUR. To the cutter of the cut of the

CONTAINER HANDLING: If empty - Nonrefillable container. Do not reuse or refill this container. Place in trash or offer for recycling if available. Do not contaminate feed or foodstuffs by storage or disposal.

PESTICUES STORAGE. Store in code, dry place.

PESTICUES CORPOSAL. IF parily filled - Call your local solid waste agency for disposal instructions.

Never place unused product down any indoor or outdoor drain.

JA2092IO ONA 3DAROTZ

HOWE REMOURS FOR USE OF UNDER STATE OR OTHER COMMERCIAL USE.

WORT FOR USE OF LANT SERING ENGINE TO BE SALE OR OTHER COMMERCIAL USE.
Weshing household lake or household with a case and work backward to avoid contact with dusted surfaces. Dust lightly to cover both upper and lower leaf surfaces with a thin, even film of that. (Many Insects bride on the underside of the leaf).

Between the surfaces with a thin even film of that. (Many Insects bride on the underside of the leaf).

Between the surfaces with a thin even film of the transmission, preferably before damage from feeding occurs. Mote the pre-harvest (time between use and harm part of the part of th

HOME GARDENS: FOR USE ON HOME GARDENS.

To control Heas, Ticks and Lice on Dogs.

To control Heas, Ticks and Lice on Dogs.

Desproach underlied in a week a see. Wearing household latex or rubber gloves dust animals thoroughly, working the dust in pit his his down to the skin. Pay close stending underline pets ayes, nose, mouth, and genital areas. Close stending beginning the less of the Lice of the Company of the Comp

POUČTRY: To control Northern Fowl Mites, apply at a rate of 1 lb. per 100 birds. Ensure thorough treatment of vent area. Lice or Swine Direct Application: Apply only 1 ounce per head as a uniform coat to the head, shoulder and back by use of a shaker can or suitable mechanical dust applicator.
Repeat as renessary, but not more often than once every 10 days. In severe infestation, both animals and the bedding may be treated. But not ship animals for slaughter within 5 days of treatment.

HORSES: Dust horse for Horn flies with shaker can or dust glove along back, neck and legs.

Horn Files, Lice Direct Application: Apply 2 oz. of dust per animal by shaker can over the head, neck, shoulders and back and tailhead. Repeat as necessary, but not more For dairy cows, bags may be suspended in the exit through which the cows leave the milking barn. The bags should hang 4 to 6 inches below the back line of the cattle. For reduction of face files, bags must be located so animals will be forced to use them also have a height so that the face is dusted.

Can be used in dust bags, shaker can and mechanical dust applicator.

Horn Files, Lice, Face Files: Place contents of this package in any commercially available dust bag, suspend bag in areas frequented by cattle or in gateways or lanes through

Horn Files, Lice, Face Files: Place contents of this package in any commercially available dust bag, suspend bag in areas frequented by cattle or in gateways or lanes through

Which the animals must pass daily for water, feed, or minerals. Bags may also be placed in loafing sheds or in front of mineral becelers. SEEF AND DAIRY CATTLE AND HORSES