

FIFRA Section 2(ee) Recommendation



FOR DISTRIBUTION AND USE ONLY IN THE STATE OF ARIZONA



EPA Reg. No. 59639-243

FOR AERIAL APPLICATIONS IN BRASSICA (COLE) LEAFY VEGETABLES

This FIFRA Section 2(ee) Recommendation Expires December 31, 2029.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This 2(ee) recommendation contains directions for use which do not appear on the package label and must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label when using *Senstar* Insecticide alone or in tank mix combinations. The most restrictive labeling applied when using a tank mix.

CROPS	PESTS	PRODUCT RATES PER ACRE	SPECIAL INSTRUCTIONS
BRASSICA (COLE) LEAFY VEGETABLES (Crop Group 5) Broccoli; Broccoli, Chinese (Gai Lon); Broccoli Raab (Rapini); Brussels Sprouts; Cabbage; Cabbage, Chinese (Bok Choy); Cabbage, Chinese (Napa); Cabbage, Chinese Mustard (Gai Choy); Cauliflower; Cavalo Broccolo; Collards; Kale; Kohlrabi; Mizuna; Mustard Greens; Mustard Spinach; Rape Greens	Aphids Banded Wing Whitefly Greenhouse Whitefly Silverleaf Whitefly Swede Midge Sweetpotato Whitefly <i>Suppression only</i> Diamondback Moth Onion Thrips (larvae)	6 to 10 fl oz (0.047 to 0.078 lb ai/A spirotetramat) (0.016 to 0.027 lb ai/A pyriproxyfen)	Apply with air equipment in adequate water for uniform coverage (minimum of 5 gallons/A). Higher water volumes may provide improved insect control. Begin application when whiteflies reach threshold levels per University/Extension recommendations (e.g., 3 to 5 adults per leaf and/or equivalent levels of immatures). Only whole fields should be treated.

Restrictions

- Do not apply more than 10 fl oz of Senstar Insecticide (0.078 lb spirotetramat and 0.027 lb pyriproxyfen) per acre per application.
- Do not make more than 2 applications per crop season.
- Do not apply more than 20 fl oz Senstar Insecticide (0.156 lb spirotetramat and 0.055 lb pyriproxyfen) per acre per crop season.
- Do not apply within 7 days of harvest.
- Minimum interval between applications is 14 days.
- Regardless of formulation do not apply more than 0.16 lb spirotetramat and 0.134 lb pyriproxyfen per acre per crop season.
- Certain nonionic and organosilicone adjuvants, which may potentially be used with Senstar Insecticide, have caused intolerable damage to bok choy, napa, mustard spinach, mizuna, and mustard greens when applied alone, in the absence of Senstar Insecticide. Testing has shown that Senstar Insecticide does not increase the potential for damage when used in combination with such adjuvants. Due to the wide variety and composition of spray adjuvants that may be used in combination with Senstar Insecticide, only use a spreading-penetrating adjuvant that is known to be safe to the target crop.
- Do not apply during bloom on crops grown for seed production.

Please contact Valent U.S.A. LLC at 1-800-6-VALENT (682-5368) to determine if this product is registered in your state.

©2024 Valent U.S.A. LLC

Registrant: Valent U.S.A. LLC
P.O. Box 5075
San Ramon, CA 94583