

RESTRICTED USE PESTICIDE

Due to acute oral, dermal, and inhalation toxicity and avian hazards
For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.



FIFRA 2(ee)
Use Directions For Application in
Soybean for Suppression of Soybean
Gall Midge

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATES OF IOWA, KANSAS,
MINNESOTA, MISSOURI, NEBRASKA, NORTH DAKOTA AND SOUTH DAKOTA

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the Environmental Protection Agency (EPA)

THIMET[®] 20-G Soil and Systemic Insecticide

EPA Reg. No.: 5481-530

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

THIS BULLETIN MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF APPLICATION.

Before applying Thimet 20-G Soil and Systemic Insecticide please read the complete label affixed to the product container. All applicable directions, restrictions and precautions on the EPA-registered label are to be followed. Use of Thimet 20-G Soil and Systemic Insecticide according to this recommendation is subject to the use precautions and limitations imposed by the label on the container.

IMPORTANT: Read the entire Directions for Use and the Limited Warranty and Disclaimer on the EPA-registered Thimet 20-G Soil and Systemic Insecticide label before using this product. Thimet 20-G is available for use only in SmartBox[®] and Lock'N Load[®] Closed Loading Systems.

Use Information

Thimet 20-G Soil and Systemic Insecticide may be applied at planting in soybean for suppression of Soybean Gall Midge larvae which feed internally within soybean stems.

Crop	Pest	Thimet 20-G Rate	Application
Soybean	Soybean Gall Midge Larvae	9.0 oz per 1000 row ft on 30-inch minimum row spacing or no more than 9.8 lb/A	Apply granules in a T-band over the row directly behind the planter shoe and in front of the press wheel, ensuring granules are lightly incorporated into soil.
<p><u>Herbicide Use Precautions:</u> Caution is recommended with use of Thimet 20-G at planting on coarse soil textures with organic matter less than 1.5% in conjunction with either preplant or preemergence applied flumioxazin-, saflufenacil- or chlorimuron-containing herbicides. Metribuzin-containing herbicides applied prior to or following Thimet 20-G applied at planting may result in crop injury. For herbicide use in conjunction with Thimet 20-G, consult the herbicide label for any crop safety precautions prior to or following use of a soil-applied systemic organophosphate insecticide.</p> <p><u>Restrictions:</u> Do not make more than one application of Thimet 20-G per year. Do not place Thimet 20-G granules in direct contact with seed. Crop Rotation: Any crop on the Thimet 20-G label can be replanted immediately. All other crops not labeled can be planted at 9 months after application.</p>			

©2022 AMVAC Chemical Corporation.

AMVAC logo, Lock’N Load, SmartBox and Thimet are trademarks owned by AMVAC Chemical Corporation.

Registrant:

AMVAC Chemical Corporation
4695 MacArthur Court, Suite 1200
Newport Beach, CA U.S.A.
1-888-462-6822